



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁴ : C07K 13/00, C12N 15/00 C12P 21/02, A61K 37/02 // A61K 35/50</p>	<p>A1</p>	<p>(11) International Publication Number: WO 89/ 08667</p> <p>(43) International Publication Date: 21 September 1989 (21.09.89)</p>
<p>(21) International Application Number: PCT/NL89/00008</p> <p>(22) International Filing Date: 10 March 1989 (10.03.89)</p> <p>(31) Priority Application Number: 1319/88</p> <p>(32) Priority Date: 11 March 1988 (11.03.88)</p> <p>(33) Priority Country: DK</p> <p>(71) Applicant (for all designated States except US): ERASMUS UNIVERSITEIT [NL/NL]; P.O.B. 1738, NL-3000 DR Rotterdam (NL).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only) : DROP, Stenwert, Leonard, Sebastian [NL/NL]; Leeuwerik Laan 12, NL-3055 EB Rotterdam (NL). BRINKMAN, Arend [NL/NL]; Van der Heimstraat 20, NL-3135 NL Vlaardingen (NL).</p>	<p>(74) Agent: OCTROOIBUREAU VRIESENDORP & GA-ADE; P.O. Box 266, NL-2501 AW The Hague (NL).</p> <p>(81) Designated States: AT, AT (European patent), AU, BB, BE (European patent), BG, BJ (OAPI patent), BR, CF (OAPI patent), CG (OAPI patent), CH, CH (European patent), CM (OAPI patent), DE, DE (European patent), DK, FI, FR (European patent), GA (OAPI patent), GB, GB (European patent), HU, IT (European patent), JP, KP, KR, LK, LU, LU (European patent), MC, MG, ML (OAPI patent), MR (OAPI patent), MW, NL, NL (European patent), NO, RO, SD, SE, SE (European patent), SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI patent), US.</p> <p>Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: RECOMBINANT IGF BINDING PROTEIN (IBP-I)</p>		
<p>(I)</p>		
<p>A-P-W-Q-C-A-P-C-S-A-E-K-L-A-L-C-P-P-V-S-A-S-C-S-E-V-T-R-S-A- G-C-G-C-C-P-M-C-A-L-P-L-G-A-A-C-G-V-A-T-A-R-C-A-R-G-L-S-C-R- A-L-P-G-E-Q-Q-P-L-H-A-L-T-R-G-Q-G-A-C-V-Q-E-S-D-A-S-A-P-H-A- A-E-A-G-S-P-E-S-P-E-S-T-E-I-T-E-E-E-L-L-D-N-F-H-L-M-A-P-S-E- E-D-H-S-I-L-W-D-A-I-S-T-Y-D-G-S-K-A-L-H-V-T-N-I-K-K-W-E-P-C- R-I-E-L-Y-R-V-V-E-S-L-A-K-A-Q-E-T-S-G-E-E-I-S-K-F-Y-L-P-N-C- N-K-N-G-F-Y-H-S-R-Q-C-E-T-S-M-D-G-E-A-G-L-C-W-C-V-Y-P-W-N-G- K-R-I-P-G-S-P-E-I-R-G-D-P-N-C-Q-M-Y-F-N-V-Q-N-</p>		
<p>(57) Abstract</p> <p>An IGF binding protein which has the amino acid sequence (I), or an equivalent modification thereof, such as a glycosylated modification. Further is indicated a DNA-sequence, coding for the protein, an expression vector and a pharmaceutical preparation containing the protein. The protein is effective as a potentiator for the function of IGF-compounds.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

Recombinant IGF binding protein (IBP-1)

5 The present invention relates to an Insulin-like Growth Factor Binding Protein, IBP-1, having a molecular weight of about 28 kD, derived from human placenta/endometrium, and equivalent modifications thereof.

10 The invention further relates to a DNA-structure, coding for the IBP-1, expression vectors containing this DNA-structure and procaryotic or eucaryotic cells comprising such a vector.

The invention still further relates to pharmaceutical preparations comprising IBP-1.

15 IGF or Insulin like Growth Factor is synonymous with somatomedins. The family of somatomedins are members of a group of polypeptides derived from the insulin gene. The gene products include insulin, insuline-like growth factor (IGF)I and II, relaxin and the B-unit of nerve growth factor (NGF) (Rinderknecht and Humbel, 1978a; Rinderknecht and Humbel, 1978b; Bradshaw 1978; 20 Isaacs et al., 1978). IGF elicits classical insulin effects on all target tissues of insulin, i.e. IGF increases glucose metabolism of adipose tissue and stimulates lipid, glycogen, and protein synthesis. IGF also stimulates DNA synthesis in certain cell types. 25 This feature reflects the capacity of IGFs to induce cell proliferation and promote organ growth in vivo. Furthermore, IGF acts on differentiation of mesenchymal cells (Froesch et al., 1985).

30 IGF-I and IGF-II are unique in that they are complexed to specific binding proteins in plasma (Smith, 1984). At least two different binding proteins have been iden-

tified in adult human serum, namely (1) binding protein 53 (BP-53) a GH dependent binding protein, believed to be derived from the 150 kD complex which carries most of the endogenous IGF peptides, (2) IBP-1, an IGF binding protein of about 30-40 kD which is tissue specifically expressed in endometrium and liver and accounts for most of the unsaturable binding sites in plasma. While the 53 kD-binding protein is under GH control the 30-40 kD species appears to be expressed in a GH independent way.

The lower molecular weight binding protein was initially identified in human amniotic fluid and has been purified and characterized (Chochinov et al., 1977; Drop et al., 1979; Drop et al., 1982). This 30-40 kD IGF binding protein appears to be identical to binding proteins that have been purified from human serum and the human hepatoma cell line, HEPG2 (Drop et al., 1984a; Pova et al., 1984; Pova et al., 1985). Pova et al showed that the NH₂-terminal amino acid sequence of the binding protein found in amniotic fluid and from the HEPG2 cell line are similar (Pova et al., 1985).

Placental protein PPl2, a protein originally isolated from human placenta, was found to bind IGF as well as to have an identical NH₂-terminal amino acid sequence (Koistinen et al., 1986).

As to the biological function of IGF-binding protein both stimulatory and inhibitory effects have been described.

Stimulatory effects of IGF-binding protein has been shown in at least two cases. Clemmons et al (1986) showed increased binding to fibroblast and smooth muscle cell surface receptors of IGF in complex with its binding protein.

Inhibitory effects of IGF-binding protein on various IGF actions in vitro, including stimulation of glucose transport by adipocytes, sulphate incorporation by chondrocytes and thymidine incorporation in fibroblasts have been described (Zapf et al., 1979; Drop et al., 1979; Ooi et al., 1986). In addition, inhibitory effects of IGF-binding proteins on growth factor mediated mitogen activity in normal cells (cartilage assay, Drop, thesis, 1983).

According to the invention, the IGF-binding protein has the following amino acid sequence:

Ala-Pro-Trp-Gln-Cys-Ala-Pro-Cys-Ser-Ala-Glu-Lys-Leu-Ala-Leu-Cys-Pro-Pro-Val-Ser-Ala-Ser-Cys-Ser-Glu-Val-Thr-Arg-Ser-Ala-Gly-Cys-Gly-Cys-Cys-Pro-Met-Cys-Ala-Leu-Pro-Leu-Gly-Ala-Ala-Cys-Gly-Val-Ala-Thr-Ala-Arg-Cys-Ala-Arg-Gly-Leu-Ser-Cys-Arg-Ala-Leu-Pro-Gly-Glu-Gln-Gln-Pro-Leu-His-Ala-Leu-Thr-Arg-Gly-Gln-Gly-Ala-Cys-Val-Gln-Glu-Ser-Asp-Ala-Ser-Ala-Pro-His-Ala-Ala-Glu-Ala-Gly-Ser-Pro-Glu-Ser-Pro-Glu-Ser-Thr-Glu-Ile-Thr-Glu-Glu-Glu-Leu-Leu-Asp-Asn-Phe-His-Leu-Met-Ala-Pro-Ser-Glu-Glu-Asp-His-Ser-Ile-Leu-Trp-Asp-Ala-Ile-Ser-Thr-Tyr-Asp-Gly-Ser-Lys-Ala-Leu-His-Val-Thr-Asn-Ile-Lys-Lys-Trp-Lys-Glu-Pro-Cys-Arg-Ile-Glu-Leu-Tyr-Arg-Val-Val-Glu-Ser-Leu-Ala-Lys-Ala-Gln-Glu-Thr-Ser-Gly-Glu-Glu-Ile-Ser-Lys-Phe-Tyr-Leu-Pro-Asn-Cys-Asn-Lys-Asn-Gly-Phe-Tyr-His-Ser-Arg-Gln-Cys-Glu-Thr-Ser-Met-Asp-Gly-Glu-Ala-Gly-Leu-Cys-Trp-Cys-Val-Tyr-Phe-Trp-Asn-Gly-Lys-Arg-Ile-Pro-Gly-Ser-Pro-Glu-Ile-Arg-Gly-Asp-Pro-Asn-Cys-Gln-Met-Tyr-Phe-Asn-Val-Gln-Asn

The complete nucleotide sequence of the corresponding cDNA sequence was shown to have the structure depicted in claim 5.

The protein according to the invention may be used as an effective potentiator for the functioning of

- somatomedins. The effect can be mediated through the firm binding between the somatomedins and their binding proteins under physiological conditions. Such complexed somatomedins together with their binding proteins are
- 5 protected against undue proteolysis, causing a significant increase of the biological half life of somatomedins. Furthermore, this IGF-binding protein or modifications hereof might function as a potent carrier of IGF to its local sites of action.
- 10 The studies of Clemmons (1986) demonstrate the potential usefulness of this IGF-1 binding protein or modifications hereof in fertilization and in potentiation of growth of connective tissue and muscle cells in tissue repair.
- As IGF-binding protein, or modifications thereof, such
- 15 as alpha 1 PEG, are the major secretory soluble protein of decidual cells of the endometrium, IBP-1 may have an important function in restricting trophoblast invasion into the endometrium during placental development. Furthermore, the inhibitory function of IBP-1 in cellular
- 20 proliferation assays and the unexpected direct inhibitory effect of IBP-1 on the oestrogen response on certain cancer cells make IBP-1 or modifications hereof a potential anticancer reagent with local growth inhibitory effect.
- 25 The invention is explained more in detail in the following description with reference to the drawing, in which
- fig. 1 illustrates a restriction map and sequence strategy for human 28 kD IGF binding protein cDNA clones,
- fig. 2 shows the nucleotide and deduced amino acid
- 30 sequence of human IGF binding protein where differences between the placental cDNA sequence and the liver cDNA sequence are shown in parenthesis, and

fig. 3 represents an SDS/PAGE analysis of culture media of COS-1 cells transfected with pSV19, pSV4, pSV4Inv and untransfected COS-1 cells.

5 The cDNA encoding IGF-BP was obtained by screening a human placental and a human hepatoma (HEPG-2) cDNA-expression library with a polyclonal antibody to human amniotic fluid binding protein (Drop et al., 1984a).

10 Restriction analysis indicated that the clones isolated from the placenta library and the clones isolated from the HEPG2 library were colinear (fig. 1), supporting their candidacy as IGF binding protein clones. The composite restriction map is shown at the top of the figure 1. A putative leader sequence is shown in front of the open box representing translated regions. Each
15 arrow shows the direction and extend of sequencing by chain termination. Four different clones are aligned. p4 and p19 originate from the placental cDNA library, while w61 and w85 originate from the HEPG2 cDNA library, (E = EcoRI, P = PstI, B = BamHI, H = HindII, S = SstI,
20 X = XbaI, N = NcoI).

The complete nucleotide sequence of the cDNA insert of one of the clones isolated (p19) was determined. The 1421 nucleotide sequence shown in fig. 2 contains a 5' untranslated region of 52 nucleotides followed
25 by an ATG codon and an open reading frame of 776 nucleotides. The potential initiation codon is flanked by 5 sequences matching Kozak's criteria for an initiation codon (Kozak, 1986). At the 3' end the open reading frame is flanked by a translation termination codon
30 (TGA) and a 569 nucleotides long 3' untranslated sequence.

The open reading frame in cDNA clone p19 has a coding capacity for a protein of 259 residues also shown in

figure 2 (by the one-letter code), with a calculated
Mw of 28,172 daltons. The initiation methione is the
first amino acid of a 24-residue highly hydrophobic
peptide sequence (underlined), representing the sequence
5 of a putative signal sequence necessary for transfer
of the nascent polypeptide sequence into the membranes
of the endoplasmatic reticulim. A favourable signal
peptidase cleavage site (ala-gly) occurs immediately
N-terminally of the alanine residue at pos +1 (von
10 Heijne, 1987). The NH₂-terminus of the predicted mature
protein is identical to the chemically determined NH₂-
terminus described for the IGF-binding protein isolated
both from amniotic fluid (Povoa et al., 1984), and
from the HEPG2 cell line (Povoa et al., 1985) and from
15 serum (Baxter et al., 1987).

Omitting the signal peptide sequence, the M_r of this
gene product is predicted to be 2,350 daltons. The
M_r of serum IGF binding protein is about 28,000 daltons
(Baxter et al., 1987). The difference is believed to
20 be accounted for by glycosylation of the IGF binding
protein (Bohn et al., 1980; Koistinen et al., 1986).

The amino acid sequence did not disclose N-linked glyco-
sylation sites (N-T, N-S). However, at least five po-
tential O-linked glycosylation sites were found in
25 the NH₂-terminal of the molecule. A RGD sequence in
the COOH terminal part of the IBP-1 protein has been
found. Such a short sequence is considered to be im-
portant for cellular attachment of matrix proteins,
such as fibronectin, vitronectin and von Willebrand
30 factor, to receptors of the integrin family.

Amino acid homology with other known proteins and pep-
tides were determined by searching the NFBR data base,
version 12.0 and 26.0. The IGF-BP protein did not show

any significant protein homology, indicating that IGF-BP is a unique protein. In particular, comparison of amino acid sequence of IGF-I, IGF-II, Insulin, Type I IGF receptor and the type II IGF receptor amino acid sequences revealed no homologous domains. In addition, no homology was found with the reported NH₂-terminal amino acid sequence for the high molecular weight IGF binding protein (Baxter et al., 1986).

It was further possible to express IBP-1 in mammalian cells. Expression vectors, pSV19, pSV4 and pSV4Inv were constructed by inserting the full length clones p4 and p19 in the expression vector pSV328. The vectors pSV4, pSV19 and pSV4Inv, in which the cDNA insert is in 3' - 5' orientation, were transfected to COS-1 cells. By use of SDS/PAGE analysis cell culture media from COS-1 cells transfected with pSV19 (lane A), pSV4 (lane B), and amniotic fluid (lane C) were analysed. The IGF binding proteins were made visible by immuno staining as described for the screening of the cDNA libraries. In culture media of pSV4 and pSV19 transfected COS-1 cells in which the gene is in the correct orientation a protein of 32 kD being immunologically indistinguishable from the IGF binding protein from amniotic fluid (fig. 3) was detected. In all culture media a band was visible which reacted with the 35 kD SMBP antibody but which was absent in the culture medium from untransfected COS-1 cells.

Although IBP-1 successfully has been expressed in COS-1 cells the lack of N-linked glycosylation sites in the putative protein also favour expression in yeast and bacteria to increase the IBP-1 production to be used in a variety of therapeutic compositions.

The invention provides therapeutic compositions comprising IBP-1 or derivatives thereof and pharmacologically

acceptable excipients. Such compositions including the IGF-binding protein or derivatives hereof according to this invention have many therapeutic uses involving the physiological functions of somatomedins.

5 The IGF-binding protein of the invention may be formulated as pharmaceutical preparations comprising the IGF-binding protein of the invention together with the usual excipients. Pharmaceutical preparations according to the invention may be in the form of suspension
10 or solutions for parenteral administration, e.g. i.v., s.c., i.m., implants, subcutaneous or intervenous administration or administration through the mucosa, e.g. oral, nasal, buccal, sublingual or rectal administration or transdermal administration.

15 For example in cases where somatomedins have to be transported to specific target tissues in a way where the physiological halflife of the somatomedins has to be increased by complexing IBP-1 described in this invention to IGF-1 and IGF-2. In accordance with this
20 invention a slow release of active IGF-1 or 2 from such complexes would ascertain a constant level of somatomedins either locally or systemically dependent upon the way of administration. IBP-1 describes in this invention hereby abolishes the potent mitogenic
25 effect of the somatomedins that administered in high dosis, i.e. intra venously, would cause unwanted local cellular proliferations in a variety of cells like fibroblasts, muscle cells and endothelial cells.

However, the IBP-1 described by this invention administered together with IGF-1, IGF-2 and other growth factors or formulated as common preparations for topical
30 use (such as PDGF, EGF, FGF, TGFalpha or TGFbeta) employed in therapeutical devices to be used in healing of wounds or in treatment of osteoporosis and in healing

of bones might be valuable for a steady and controlled release of the somatomedins in such therapeutical devices.

Such preparations may optionally be administered in the form of combination preparations e.g. comprising IBP-1 and IGF-1, IBP-1 and IGF-2 or IBP-1, IGF-1 and IGF-2.

In general, IBP-1 according to this invention might turn out to exhibit a potent regulatory function in the release of IGF-1 and/or IGF-2 in future treatment of injuries or other malfunctions that requires increased IGF-1 and/or IGF-1 levels.

On the other hand IBP-1 or derivatives thereof according to this invention might be useful in therapy of the proliferation of certain cancers characterized by producing somatomedins in high amounts thus inhibiting the autocrine/paracrine physiological stimulation of unwanted cellular proliferation in cancers like chondrosarcomas, fibrosarcomas, and mammacarcinomas.

Furthermore, the IBP-1 or derivatives hereof described in this invention is useful for the production of antibodies. Such mono- or polyclonal antibodies are suitable for developing immunological methods like immunohistochemical analysis of IBP-1 in tissues and for developing ELISA for IBP-1 quantitation. Such ELISA will prove valuable for early screening the levels of IBP-1 in patients with altered IGF-1 and 2 levels.

Pharmaceutical preparations of this invention for s.c. and i.m. administration can be prepared by mixing the following constituents: IBP-1 and derivatives thereof together with IGF-1, IGF-2 and other growth factors, an isotonic agent, a buffer, a preservative and water.

After mixing the pH value of the preparation is, if necessary, adjusted to pH = 7.3.

Examples of preservatives: phenol and m-cresol. Examples of an isotonic agent: sodium chloride and glycerol.

5 Example of buffer is sodium phosphate.

Pharmaceutical preparations of this invention for trans-mucosal administration can be prepared by mixing the following constituents: IBP-1 and derivatives thereof, together with IGF-1, IGF-2 and other growth factors,
10 a buffer, an isotonic agent, a preservative, an absorption promotor and a vehicle e.g. water, cellulose, water-soluble cellulose alkylethers, crystalline cellulose, water-soluble polyacrylates or mixtures thereof.

Pharmaceutical preparations of this invention for trans-dermal administration can be prepared by mixing the following constituents: IBP-1 and derivatives thereof together with IGF-1, IGF-2 and other growth factors,
15 an isotonic agent, a preservative and a vehicle e.g. a hydrophilic gel of water-soluble cellulose alkyl-ethers.
20

This invention is further explained in the following working example describing the isolation and characterization of IBP-1.

EXAMPLE

25 Screening the Lambda gt11 Expression Library

A human placenta cDNA library in lambda gt11 and a cDNA library of the human hepatoma cell line HEPG2 were screened with a polyclonal antibody to human amniotic fluid binding protein according to the procedure
30 described by Young and Davis (Young and Davis, 1982).

Rabbit antibody to 35 kD somatomedin binding protein SMBP isolated from human amniotic fluid was produced and purified as described by Drop et al., 1984a. The antibody was absorbed against E.coli Y1090 and lambda gt11 proteins by incubating with nitrocellulose filters that had been lifted from confluent lysis plates of E.coli Y1090/lambda gt11 induced with 10 mM isopropyl beta-d-thiogalacopyranoside (IPTG). The antibody was further absorbed against human serum albumin immobilized on nitrocellulose filters. Approximately 4×10^5 clones of the placental library were screened and about 0.5×10^5 of the HPEG2 library. $3-5 \times 10^4$ plaque forming units per 150 mm Petri dish were plated on a lawn of Y1090 bacteria and incubated. After 2 hr incubation the plates were covered with nitrocellulose filters (Millipore HATF) that had been saturated with 10 mM IPTG. The plates were incubated at 37°C for 2-2.5 hrs. The filters were removed, washed with Tris-buffered saline (TBS; 10 mM Tris/HCl, pH 7.5/150 mM NaCl) at room temperature and incubated with 3% BSA in TBS for 30 min. at room temperature. Partly purified rabbit polyclonal 35 kD SMBP antibody diluted 1:125 was added to 3% BSA in TBS plus 0.02% azide, and the filters were incubated overnight at 4°C. The filters were washed and incubated for 60 min. at room temperature with horse-radish peroxidase conjugated goat anti-rabbit IgG (Tago) diluted 1:200 in 3% BSA in TBS. The filters were washed and stained with amidophenyl and naphthol AS-MX phosphate in 0.2 M Tris/HCl, pH 9.2, 10 mM $MgCl_2$ at room temperature.

Positive phages were isolated and DNA was isolated by standard methods (Maniatis et al., 1982). About 33 plaques strongly cross-reacting with the polyclonal antibody were identified in the placenta and HPEG2 cDNA library. Following re-screening inserts varying in size between 0.9-1.5 Kb were isolated and subcloned

in the vector PTZ19 from Pharmacia. All isolated clones showed cross-hybridization in a Southern blot except one clone from the placenta library and the 5 weakly hybridizing clones from the HEPG2 library.

- 5 DNA was digested with various restriction endonucleases (BRL, NEN, Boehringer) according to the suppliers directions, electrophoresed in 0.8% agarose, and transferred to nitrocellulose filters according to the method of southern (Southern, 1975). mRNA was denaturated with
10 dimethylsulfozide (DMSO) and glyoxal, subjected to electrophoresis in 1% agarose and transferred to nitrocellulose filters (Millipore HFTF).

- Restriction fragments were subcloned in the vectors PTZ18 or PTZ19 (Pharmacia) and sequenced according
15 to the chain termination method (Sanger et al., 1977). In regions which lacked convenient restriction sites, appropriate clones were generated by Bal 31 nuclease digestion.

Transfection of COS-1 Cells

- 20 The full length cDNA clones p4 and p19 were subcloned in the EcoR1 site of pSV328, which expressed cloned inserts using the simian virus 40 (SV40) early promotor (Van Heuvel et al., 1986). A DEAE-dextran procedure (McCuthchan & Pagano, 1986) followed by treatment with
25 100 μ M chloroquine in Dulbecco's MEM (DMEM) for 4 hrs was used to transfect COS-1 cells (Gluzman, 1981). After this treatment the cells were grown 24 hrs with DMEM plus 5% foetal calf serum. Medium was removed after 72 hrs, and the cells were washed extensively
30 with DMEM and incubated for 72 hrs with DMEM without serum. Production of 32 kD binding protein in culture media was determined using 35 kD SMBP antibody.

Purification of IBP-1

Proteins from amniotic fluid or from conditioned media were precipitated with ammonium sulphate at a final concentration of 35%. Following centrifugation the supernatant was brought to 50% ammonium sulphate. The pellet was dissolved in 45% ammonium sulphate and the final pellet was dissolved in 50 mM Tris HCl, pH 7.5 for further purification and characterization. The dissolved ammonium sulphate precipitate was further purified by reverse phase chromatography on C₁₈. Following washings with 50 mM Tris-HCl, pH 7.5 and Tris HCl, pH 7.5 in 50% methanol the pure IBP-1 was eluted from the column with Tris-HCl, pH 7.5 in 65% methanol. The IBP-1 was precipitated overnight in a 7% Trichloroacetic-acid solution. The precipitate was dissolved in 20 mM Tris-HCl, pH 7.5, lyophilized and stored at \pm 20°C.

Characterization of purified IBP-1

IBP-1 binds both IGF-1 and IGF-2 with approximately the same specificity when measured in binding assays with [¹²⁵I]IGF-1 and [¹²⁵I]IGF-2 with or without excess of cold IGF-1 or IGF-2. The specificity of IBP-1 reaction was tested in similar assays employing such competition assays and in assays in which IBP-1 - IGF binding was visualized through the specific reaction of antibody against IBP-1. In such assays IBP-1 did only react with IGF-1 and IGF-2 but not with closely related compounds like insulin, proinsulin or truncated forms thereof.

The biological effects of purified samples of IBP-1 were tested in in vitro mitogenicity tests employing ³H-thymidin incorporation into a MCF-7 mamma carcinoma cell line. The stimulatory effect of both IGF-1 and

IGF-2 on cell proliferation was inhibited by IBP-1 in a dose dependent way. The effect of IBP-1 on IGF-2 dependent cell proliferation was more pronounced than that of the effect of IBF-1.

- 5 Furthermore, in an assay employing ^{35}S -Methionin incorporation into cartilage the stimulatory effect of IGF-1 and IGF-2 was abolished by IBP in the low ng range.

Examples of pharmaceutical preparations:

10 EXAMPLE 1

25 ng IBP-1
10 ng IGF-1
0.7% NaCl
1/75M sodiumphosphate
15 water ad 1 ml

The calculated amounts of IBP-1 and IGF-1 were dissolved and diluted in phosphate buffer containing NaCl. The pH was adjusted to 7.3-7.4.

EXAMPLE 2

20 25 ng IBP-1
10 ng IGF-2
1.6% glycerin
1/75M sodiumphosphate
0.01% benzalconiumchlorid
25 0.05% sodiumedetat
water ad 1 ml

The calculated amounts of IBP-1 and IGF-2 were dissolved and diluted in phosphate buffer containing glycerin, benzalconiumchlorid and sodiumedetat. The pH

File Appⁿation No.: 09/348,815

**NOTICE TO COMPLY WITH REQUIREMENTS FOR PATENT APPLICATIONS CONTAINING
NUCLEOTIDE SEQUENCE AND/OR AMINO ACID SEQUENCE DISCLOSURES**

The nucleotide and/or amino acid sequence disclosure contained in this application does not comply with the requirements for such a disclosure as set forth in 37 C.F.R. 1.821 - 1.825 for the following reason(s):

- ☒ 1. This application clearly fails to comply with the requirements of 37 C.F.R. 1.821-1.825. Applicant's attention is directed to these regulations, published at 1114 OG 29, May 15, 1990 and at 55 FR 18230, May 1, 1990.
- ☐ 2. This application does not contain, as a separate part of the disclosure on paper copy, a "Sequence Listing" as required by 37 C.F.R. 1.821(c).
- ☒ 3. A copy of the "Sequence Listing" in computer readable form has not been submitted as required by 37 C.F.R. 1.821(e).
- ☐ 4. A copy of the "Sequence Listing" in computer readable form has been submitted. However, the content of the computer readable form does not comply with the requirements of 37 C.F.R. 1.822 and/or 1.823, as indicated on the attached copy of the marked -up "Raw Sequence Listing."
- ☐ 5. The computer readable form that has been filed with this application has been found to be damaged and/or unreadable as indicated on the attached CRF Diskette Problem Report. A Substitute computer readable form must be submitted as required by 37 C.F.R. 1.825(d).
- ☐ 6. The paper copy of the "Sequence Listing" is not the same as the computer readable form of the "Sequence Listing" as required by 37 C.F.R. 1.821(e).
- ☐ 7. Other: _____

Applicant Must Provide:

- ☒ An initial or substitute computer readable form (CRF) copy of the "Sequence Listing".
- ☒ An initial or substitute paper copy of the "Sequence Listing", as well as an amendment directing its entry into the specification.
- ☒ A statement that the content of the paper and computer readable copies are the same and, where applicable, include no new matter, as required by 37 C.F.R. 1.821(e) or 1.821(f) or 1.821(g) or 1.825(b) or 1.825(d).

For questions regarding compliance to these requirements, please contact:

For Rules Interpretation, call (703) 308-4216

For CRF Submission Help, call (703) 308-4212

PatentIn Software Program Support (SIRA)

Technical Assistance.....703-287-0200

To Purchase PatentIn Software.....703-306-2600

PLEASE RETURN A COPY OF THIS NOTICE WITH YOUR RESPONSE

was adjusted to 7.4

EXAMPLE 3

25 ng IBP-1
10 ng IGF-1
5 10 ng IGF-2
5% hydroxyethylcellulose
0.9% benzylalcohol
1/75M phosphate buffer
water ad 1 ml

10 The gel is prepared by mixing hydroxyethylcellulose
with the waterphase containing IBP-1, IGF-1 and IGF-2.

P a t e n t C l a i m s :

1. An IGF binding protein, comprising the following amino acid sequence:

5 A-P-W-Q-C-A-P-C-S-A-E-K-L-A-L-C-P-P-V-S-A-S-C-S-E-V-T-R-S-A-
G-C-G-C-C-P-M-C-A-L-P-L-G-A-A-C-G-V-A-T-A-R-C-A-R-G-L-S-C-R-
A-L-P-G-E-Q-Q-P-L-H-A-L-T-R-G-Q-G-A-C-V-Q-E-S-D-A-S-A-P-H-A-
A-E-A-G-S-P-E-S-P-E-S-T-E-I-T-E-E-E-L-L-D-N-F-H-L-M-A-P-S-E-
E-D-H-S-I-L-W-D-A-I-S-T-Y-D-G-S-K-A-L-H-V-T-N-I-K-K-W-E-P-C-
R-I-E-L-Y-R-V-V-E-S-L-A-K-A-Q-E-T-S-G-E-E-I-S-K-F-Y-L-P-N-C-
10 N-K-N-G-F-Y-H-S-R-Q-C-E-T-S-M-D-G-E-A-G-L-C-W-C-V-Y-P-W-N-G-
K-R-I-P-G-S-P-E-I-R-G-D-P-N-C-Q-M-Y-F-N-V-Q-N-

or an equivalent modification thereof.

2. A glycosylated modification of the protein according to claim 1.

15 3. An IGF binding protein according to claim 2, in which one or more hydroxy groups are glycosylated.

4. A DNA sequence, coding for the IGF binding protein as defined in claim 1.

20 5. A cDNA sequence according to claim 4 in which the coding strand includes the following structure:

1 GGGCGGGCAC AGCCAGAGAG CATCGGCCCC TGTCTGCTGC TCGCGCCTGG
51 AGATGTCAGA GGTCCCCGTT GCTCGCGTCT GGCTGGTACT GCTCCTGCTG
101 ACTGTCCAGG TCGGCGTGAC AGCCGGCGCT CCGTGGCAGT GCGCGCCCTG
151 CTCCGCCGAG AAGCTCGCGC TCTGCCCGCC GGTGTCCGCC TCGTGCTCGG
201 AGGTCACCCG GTCCGCCGGC TCGGGCTGTT GCCCGATGTG CGCCCTGCCT
251 CTGGGCGCCG CGTGCGGCGT GGC GACTGCA CGCTGCGCCC GGGGACTCAG
301 TTGCCGCGCG CTGCCGGGGG AGCAGCAACC TCTGCACGCC CTCACCCGCG
351 GCCAAGGCGC CTGCGTGCA GAGTCTGACG CCTCCGCTCC CCATGCTGCA
401 GAGGCAGGGA GCCCTGAAAG CCCAGAGAGC ACGGAGATAA CTGAGGAGGA
451 GCTCCTGGAT AATTTCCATC TGATGGCCCC TTCTGAAGAG GATCATTCCA
501 TCCTTTGGGA CGCCATCAGT ACCTATGATG GCTCGAAGGC TCTCCATGTC
551 ACCAACATCA AAAAATGGAA GGAGCCCTGC CGAATAGAAC TCTACAGAGT
601 CGTAGAGAGT TTAGCCAAGG CACAGGAGAC ATCAGGAGAA GAAATTTCCA
651 AATTTTACCT GCCAACTGC AACAAGAATG GATTTTATCA CAGCAGACAG
701 TGTGAGACAT CCATGGATGG AGAGGCGGGA CTCTGCTGGT GCGTCTACCC
751 TTGGAATGGG AAGAGGATCC CTGGGTCTCC AGAGATCAGG GGAGACCCCA
801 ACTGCCAGAT GTATTTTAAT GTACAAAAC GAACCAGAT GAAATAATGT
851 TCTGTCACGT GAAATATTTA AGTATATAGT ATATTTATAC TCTAGAACAT
901 GCACATTTAT ATATATGTAT ATGTATATAT ATATAGTAAC TACTTCTTAT
951 ACTCCATACA TAACTTGATA TAGAAAGCTG TTTATTTATT CACTGTAAGT
1001 TTATTTTTTC TACACAGTAA AAAC TTGTAC TATGTTAATA ACTTGTCTTA
1051 TGTCAATTTG TATATCATGA AACACTTCTC ATCATATTGT ATGTAAGTAA
1101 TTGCATTTCT GCTCTTCCAA AGCTCCTGCG TCTGTTTTTA AAGAGCATGG
1151 AAAAATACTG CCTAGAAAAT GCAAAATGAA ATAAGAGAGA GTAGTTTTTC
1201 AGCTAGTTTG AAGGAGGACG GTTAACTTGT ATATTCCACC ATTCACATTT
1251 GATGTACATG TGTAGGGAAA GTTAAAAGTG TTGATTACAT AATCAAAGCT
1301 ACCTGTGGTG ATAGTTGCCA CCTGTAAAAA TGTACACTGG ATATGTTGTT
1351 AAACACGTGT CTATAATGGA AACATTTACA ATAAATATTC TGCATGGAAA
1401 AAAAAAAAAA AAAAAAAAAA AAAAA

6. An expression vector, containing the DNA sequence defined in claim 4 or 5.
7. The expression vector pSV19.
8. The expression vector pSV4.
- 5 9. A cell line or a microorganism, comprising an expression vector in accordance with claims 6, 7 or 8.
10. A pharmaceutical preparation, comprising the binding protein defined in any of the claims 1-3.

LITERATURE:

1. Baxter R.C., Martin J.L., Tyler M.I., Howden, M.E.H.
Growth hormone-dependent insulin-like growth factor
(IGF) binding protein from human plasma differs
from other human IGF binding proteins.
a. Biochem. Biophys. Res. Commun. 1986: 139:
1256-1261.
2. Bohn H., Kraus W. Isolierung und Charakterisierung
eines neuen plaszentaspezifischen Proteins (PP12).
Arch. Gynecol. 1980: 229-279
3. Bradshaw R.A. Nerve growth factor. Annu. Rev. Bio-
chem. 1978: 47: 191-216.
4. Clemmons D.R., Elgin R.G., Han V.K.M., Casella
S.J., D'Ercole A.J., Van Wyk J.J. Cultured fibroblast
monolayers secrete a protein that alters the cellular
binding of somatomedin-C/insulin-like growth factor
I. J. Clin. Invest. 1986: 77: 1548-1556.
5. De Vroede M.A., Tseng, L.Y.H., Katsoyannis P.G.,
Nissiev S.P., Rechler M.M. Modulation of insulin-like
growth factor I binding to human fibroblast monolayer
cultures by insulin-like growth factor carrier
proteins released to the incubation media. J. Clin.
Invest. 1986: 77: 602-613.
6. Drop S.L.S., Valiquette G., Guyda H.J., Corvol
M.T., Posner B.I. Partial purification and charac-
terization of a binding protein for insulin-like
activity (ILAs) in human amniotic fluid: a possible
inhibitor of insulin-like activity. Acta. Endocrinol.
(Copenh.) 1979) 90: 505-5118.

7. Drop S., Kortleve D., Guyda H. Isolation of a somatomedin binding protein in human amniotic fluid: development of a radioimmunoassay. *Pediatr. Res.* 1982: 16: 890894.
8. Drop S.L.S., Kortleve D.J., Guyda H.J. Isolation of a somatomedin-binding protein from preterm amniotic fluid. Development of a radioimmunoassay. *J. Clin. Endocrinol. Metab.* 1984a: 59: 899-907.
9. Elgin R.G., Busby Jr. W.H., Clemmons D.R. An insulin-like growth factor (IGF) binding protein enhances the biologic response to IGF-I. *Proc. Natl. Acad. Sci. USA* 1987: 84: 3254-3258.
10. Froesch E.R. Schmid C., Schwander J., Zapf J. Actions of insulin-like growth factor. *Ann. Rev. Physiol.* 1985: 47: 443-467.
11. Heijne G. von. A new method for predicting signal sequence cleavage sites. *Nucleic. Acid. Research* 1986: 14: 714-722.
12. Isaacs N., James R., Niall H. et al. Relaxin and its structural relationship to insulin. *Nature* 1978: 271: 278-281.
13. Koistinen R., Kalkkinen N., Huhtala M.L., Seppala M., Bohn H., Rutanen E.M. Placental protein 12 is a decidual protein that binds somatomedin and has an identical N-terminal amino acid sequence with somatomedin-binding protein from human amniotic fluid. *Endocrinology* 1986: 118: 1375-1378.
14. Kozak M. Influences of mRNA secondary structure on initiation by eukaryotic ribosomes. *Proc. Natl. Acad. Sci. USA* 1986: 83: 2850-2854.

15. Maniatis T., Fritsch E.F., Sambrook J. ed. Molecular cloning: a laboratory manual. Cold Spring Harbor Laboratory 1982.
16. Pova G., Enberg G., Jörnvall H., Hall K. Isolation and characterization of a somatomedin-binding protein from mid-term human amniotic fluid.
a. Eur. J. Biochem. 1984: 144: 199-204.
17. Pova G., Isaksson M., Jörnvall H., Hall K. The somatomedin-binding protein isolated from a human hepatoma cell line is identical to the human amniotic fluid somatomedin-binding protein. Biochem. Biophys. Res. Commun. 1985: 128: 1071-1078.
18. Rinderknecht E., Humbel R.E. The amino acid sequence of human insulin-like growth factor I and its structural homology with proinsulin. J. Biol. Chem. 1978a: 253: 2769-2776.
19. Rinderknecht E., Humbel R.E. Primary structure of human insulin-like growth factor II. FEBS Lett. 1978b: 89: 283-286.
20. Sanger F., Nicklan A.R., Coulson J. DNA sequencing with chain-terminating inhibitor. Proc. Natl. Acad. Sci. 1977: 74: 5463.
21. Smith G.L. Somatomedin carrier proteins. Mol. Cell. Endocrinal. 1984: 34: 85-89.
22. Southern E. Detection of specific sequences among DNA fragments separated by gelelectrophoresis. J. Mol. Biol. 1975: 98: 503.

23. Young R.A., Davis R.W. Efficient isolation of genes by using antibody probes. Proc. Natl. Acad. Sci. USA 1983: 80: 1194-1198.
24. Gluzman Y., SV40-transformed simian cells support the replication of early SV40 mutants, Cell., 23 (1) 175-82, 1981, Jan/IMD-8108.
25. McCutchan J.H., Pagano J.S., Enhancement of the infectivity of simian virus 40 deoxyribonucleic acid with diethylaminoethyl-dextran, J. Natl. Cancer Inst., 41 (2) 351-7, Aug. 68/IMD-6810.
26. Van Heuvel M., Bosveld I.J., Mooren A.A., Trapman J., Zwarthoff E.C., Properties of natural and hybrid murine alpha interferons, J. Gen. Virol., 67 (Pt 10) 2215-22, 1986 Oct/IMD-8701.
27. Zapf j., Schoenle E., Jagers G., Sand I., Grünwald J., Froesch E.R., Inhibition of the action of non-suppressible insulin-like activity on isolated fat cells by binding to its carrier protein. J. Clin. Invest. 1979: 63, 1077-1084.
28. Baxter R.C., Martin J.L., Binding proteins for insulin-like growth factors in adult rat serum. Comparison with other human and rat binding proteins. Biochem. Biophys. Res. Commun. 1987: 147, 408.
29. Ooi G.T., Herington A.C., Covalent cross-linking of insulin-like growth factor I to a specific inhibitor from human serum, Biochem. Biophys. Res. Commun. 1986: 137, 411.
30. Chochinov R.H., mariz I.K., Hajek A.S., Daughaday W.H., Characterization of a protein in mid-term

human amniotic fluid which reacts in somatomedin-C
radioreceptor assay, J. Clin. Endocrinol. Metab.
1977:44, 902.




FIG.1

2/3

GGGCGGGCACAGCCAGAGAGCATCGGCCCTGTCTGCTGCTCGCGCCTGGAGCTGTCAGAGGTCCCCGTTGCTCGCGTCTGGCTGGTACT -13
10 20 30 40 50 60 70 80 90

⁻²⁵ M S E V P V A R V W L V L
L L L T V Q V G V T A G ⁺¹ A P W Q C A P C S A E K L A L C P P 18
GCTCCTGCTGACTGTCCAGGTGCGCGTGACAGCGCGCTCCGTGGCAGTGCGCGCCCTGCTCCGCCGAGAAGCTCGCGCTCTGCCCGCC
100 110 120 130 140 150 160 170 180

V S A S C S E V T R S A G C G C C P M C A L P L G A A C G V 48
GGTGTCCGCTCGTCTCGGAGGTACCCCGTCCGCGGCTGCGGCTGTGCCCCGATGTGCGCCCTGCTCTGGCGCGCGCTGCGGCGT
190 200 210 220 230 240 250 260 270

A T A R C A R G L S C R A L P G E Q Q P L H A L T R G Q G A 78
GGCGACTGCACGCTGCGCGCGGGGACTCAGTTGCCGCGCGTGGCGGGGAGCAGCAACCTCTGCACGCCCTACCCCGCGCAAGCGC
280 290 300 310 320 330 340 350 360

C V Q E S D A S A P H A A E A G S P E S P E S T E I T E E E 108
CTGCGTGCAGGAGTCTGACGCTCCGCTCCCATGTCTGCAGAGGAGGAGCCCTGAAAGCCAGAGAGACGGAGATAACTGAGGAGGA
370 380 390 400 410 420 430 440 450

L L D N F H L M A P S E E D H S I L W D A I S T Y D G S K A 138
GCTCCTGGATAATTTCCATCTGATGCCCCCTTCTGAAGAGGATCATTCATCCTTTGGGACGCCATCAGTACCTATGATGGCTCGAAGGC
460 470 480 490 500 510 520 530 540

L H V T N I K K W K E P C R I E L Y R V V E S L A K A Q E T 168
TCTCCATGTCAACCAATCAAAAAATGGAAGGAGCCCTGCCGAATAGAACTCTACAGAGTCTGAGAGATTAGCCAAGGCACAGGAGAC
550 (C) 570 580 590 600 610 620 630

S G E E I S K F Y L P N C N K N G F Y H S R Q C E T S M D G 198
ATCAGGAGAAGAAATTTCCAAATTTACCTGCCAACTGCAACAAGATGGATTTTATCAGCAGACAGTGTGAGACATCCATGGATGG
640 650 660 670 680 690 700 710 720

E A G L C W C V Y P W N G K R I P G S P E I R G D P N C Q M 228
AGAGCGGGGACTCTGCTGGTGCCTTACCTTGGAAATGGGAAGAGGATCCCTGGGTCTCCAGAGATCAGGGGAGACCCCACTGCCAGAT
730 740 750 760 770 780 790 800 810

Y F N V Q N 234
GTATTTTAATGTACAAAACCTGAAACCAGATGAAATAATGTTCTGTACGTGAAATTTTAAAGTATATAGTATTTTACTCTAGAACAT
820 830 840 850 860 870 880 890 900

GCACATTTTATATATGTATATGTATATATATATAGTAACCTACTTCTTATACTCCATACATAACTTGATATAGAAAGCTGTTTTTATT 910 920 930 940 950 960 970 980 990

CACTGTAAGTTTATTTTTCTACACAGTAAAACTTGTAATTAACCTTGCTCTATGTCAATTTGTATATCATGAAACACTTCTC 1000 1010 1020 1030 1040 1050 1060 1070 (T)1080

ATCATATTGTATGTAAGTAATTGCATTCTGCTCTTCCAAAOCTCCTGCGTCTGTTTTTAAAGAGCATGAAAAAATACTGCCTAGAAAAAT 1090 1100 1110 (G) 1130 1140 (C) 1150 1160 1170

GCAAAATGAAATAAGAGAGAGTAGTTTTTCAGCTAGTTTGAAGGAGGACGGTTAACTTGATATTCCACCATTACATTTGATGTACATG 1180 1190 1200 1210 1220 1230 1240 1250 1260

TGTAGGGAAGTTAAAGTGTGATTACATAATCAAAGCTACCTGTGGTGATAGTTGCCACCTGTTAAATGTACACTGGATATGTTGTT 1270 1280 1290 1300 1310 1320 1330 1340 1350

AAACAGTGTCTATAATGAAACCTTACATAAATATTCTGCATGGAAAAAATAAAAAAAAAAAAAAAAAAAAAA 1360 1370 1380 1390 1400 1410 1420

FIG. 2

3/3





FIG. 3

INTERNATIONAL SEARCH REPORT

International Application No PCT/NL 89/00008

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) *		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC ⁴ : C 07 K 13/00, C 12 N 15/00, C 12 P 21/02, A 61 K 37/02, //A 61 K 35/50		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
IPC ⁴	C 07 K; C 12 N; C 12 P; A 61 K	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched *		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁸		
Category *	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	EP, A, 0141326 (BEHRINGWERKE) 15 May 1985, see the whole document --	1-3,10
X	Biochemical and Biophysical Research Communications, vol. 141, no. 1, 23 November 1986, Academic Press, Inc. (US) M.L. Huhtala et al.: "Biologically active domain in somatostatin-binding protein", pages 263-270 see the whole article --	1-3,10
P,X	Molecular Endocrinology, vol. 2, no. 5, May 1988 The Endocrine Society Y.L. Lee et al.: "Insulin-like growth factor (IGF) binding protein complementary deoxyribonucleic acid from human HEP G2 hepatoma cells: predicted protein sequence suggests an IGF binding domain different from those of the IGF-I and IGF-II receptors" pages 404-411 see the whole article --	1-10
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>* Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"G" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
20th June 1989		11. 07. 89
International Searching Authority		Signature of Authorized Officer
EUROPEAN PATENT OFFICE		 P.C.G. VAN DER PUTTEN

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
P,X	Biochemical and Biophysical Research, vol. 152, no. 3, 16 May 1988 Academic Press, Inc. M.T. Brewer et al.: "Cloning characterization, and expression of a human insulin-like growth factor binding protein", pages 1289-1297 see the whole article --	1-10
P,X	Embo Journal, vol. 7, no. 8, 1988 IRL Press Ltd. Oxford (GB) A. Brinkman et al.: "Isolation and characterization of a cDNA encoding the low molecular weight insulin-like growth factor binding protein (IBP-1) pages 2417-2423 see the whole article --	1-10
P,X	FEBS-Lett., vol. 236, no. 2, August 1988, Elsevier Science Publishers B.V. Biomedical Division M. Julkunen et al.: "Primary structure of human insulin-like growth factor- binding protein/placental protein 12 and tissue-specific expression of its mRNA", pages 295-302 see the whole article --	1-10
P,X	Nucleic Acids Research, vol. 16, no. 17, 1988, IRL Press Ltd. Oxford (GB) U. Grundmann et al.: "Cloning of cDNA encoding human placental protein 12 (PP12):binding protein for IGF I and somatomedin" page 8711 see the whole article -----	1-10

NL 8900008
SA 27516

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

670-115000 (0.1)

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82